

Inertial modes of neutron stars with the superfluid core

Shijun Yoshida^{1*} and Umin Lee^{2†}

¹*Centro Multidisciplinar de Astrofísica – CENTRA, Departamento de Física, Instituto Superior Técnico, Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal*

²*Astronomical Institute, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan*

Typeset 2 February 2008; Received / Accepted

ABSTRACT

We investigate the modal properties of inertial modes of rotating neutron stars with the core filled with neutron and proton superfluids, taking account of entrainment effects between the superfluids. In this paper, the entrainment effects are modeled by introducing a parameter η so that no entrainment state is realized at $\eta = 0$. We find that inertial modes of rotating neutron stars with the superfluid core are split into two families, which we call ordinary fluid inertial modes (i^o -mode) and superfluid inertial modes (i^s -mode). The two superfluids in the core counter-move for the i^s -modes. For the i^o -modes, $\kappa_0 = \lim_{\Omega \rightarrow 0} \omega/\Omega$ is only weakly dependent on the entrainment parameter η , where Ω and ω are the angular frequency of rotation and the oscillation frequency observed in the corotating frame of the star, respectively. For the i^s -modes, on the other hand, $|\kappa_0|$ almost linearly increases as η increases. Avoided crossings as functions of η are therefore quite common between i^o - and i^s -modes. We find that some of the i^s -modes that are unstable against the gravitational radiation reaction at $\eta = 0$ become stable when η is larger than η_{crit} , the value of which depends on the mode. Since the radiation driven instability associated with the current multipole radiation is quite weak for the inertial modes and the mutual friction damping in the superfluid core is strong, the instability caused by the inertial modes will be easily suppressed unless the entrainment parameter η is extremely small and the mutual friction damping is sufficiently weak.

Key words: instabilities – stars: neutron – stars: oscillations – stars : rotation

1 INTRODUCTION

Since Andersson (1998) and Friedman & Morsink (1998) found that the r -modes of a rotating neutron star are unstable against gravitational radiation reaction even for infinitesimally small rotation speeds, many papers on the r -mode instability have appeared. As a result of these investigations, the importance of the instability in astrophysics has been pointed out mainly in the following two ways, that is, the instability is strong enough to limits both the spin of newly born neutron stars and that of old accreting neutron stars, and neutron stars in which the r -mode oscillation is excited will be a potential source of gravitational waves that are detectable by the LIGO and other detectors (Lindblom, Owen, & Morsink 1998; Andersson, Kokkotas, & Schutz 1999; Owen et al. 1998; Bildsten 1998; Andersson, Kokkotas, & Stergioulas 1999). For a recent review, see, e.g., Andersson & Kokkotas (2001).

It has long been suggested that neutrons in the inner crust and neutrons and protons in the core of neutron stars are in superfluid states when the interior temperatures cool down below $T \sim 10^9$ K (e.g., Shapiro & Teukolsky 1983). Since the interior temperature of neutron stars is believed to cool down quickly by emitting neutrinos (e.g., Baym & Pethick 1979), we may expect that many observable neutron stars, except newly born ones, are likely to have a core that contains superfluids. Although one of the important ingredients that affect the oscillations of neutron stars is superfluidity in the interior, it was rather recently that Epstein (1988) pointed out the importance of superfluidity for the oscillations, suggesting the existence

* E-mail: yoshida@fisica.ist.utl.pt

† E-mail: lee@astr.tohoku.ac.jp

of a new family of acoustic modes associated with superfluidity by means of the local dispersion relations derived for the case of non-rotating neutron stars. Later on, Lindblom & Mendell (1994) studied the f -modes of rotating neutron stars that have a core filled with neutron and proton superfluids, taking account of entrainment effects between the superfluids, and showed that the ordinary fluid-like f -modes are not strongly affected by the superfluidity in the core. In the same paper, Lindblom & Mendell (1994) suggested the existence of superfluid-like acoustic modes, which they called s_0 -modes, but they could not find the s_0 -modes numerically. First numerical examples of the superfluid-like acoustic modes of neutron stars with the superfluid core were obtained by Lee (1995), who also showed that g -modes are not propagating in the core filled with superfluids. Andersson & Comer (2001) discussed the dynamics of superfluid neutron star cores examining local dispersion relations in detail, and they confirmed that g -modes are excluded from the superfluid core (Lee 1995). Applying their argument to the r -modes, they suggested that the r -modes of a superfluid neutron star core are split into two distinct families, that is, ordinary fluid-like r -modes and superfluid-like r -modes. Most recently, Prix & Rieutord (2002) have computed in great detail acoustic modes of non-rotating neutron stars composed of neutron and proton superfluids for a wide range of the entrainment parameter. Although their results are basically consistent with the results obtained by Lee (1995) and Andersson & Comer (2001), they have also shown that the assumption that $|\delta\mathbf{v}_p - \delta\mathbf{v}_n| \sim 0$ for the ordinary fluid-like acoustic modes and $|\rho_p\delta\mathbf{v}_p + \rho_n\delta\mathbf{v}_n| \sim 0$ for the superfluid-like acoustic modes is not necessarily helpful to distinguish the two families, where $\delta\mathbf{v}_p$ and $\delta\mathbf{v}_n$ are the Eulerian velocity perturbations of the proton and neutron superfluids, the mass densities of which are denoted as ρ_p and ρ_n . We also note that Sedrakian & Wasserman (2000) examined rotational effects on the fundamental modes in uniform density superfluid stars, applying the so-called tensor virial technique to normal modes. As for relativistic non-rotating superfluid stars, Comer, Langlois, & Lin (1999) calculated superfluid-like acoustic modes and showed that the basic features of the superfluid modes are quiet similar to those of the corresponding Newtonian superfluid modes. More recently, Andersson, Comer & Langlois (2002) have found avoided crossings between the superfluid- and the ordinary fluid-like acoustic modes in a relativistic superfluid neutron star as the entrainment is varied.

To give an answer to the question whether the r -mode instability will survive the damping due to mutual friction in the superfluid core of cold neutron stars, Lindblom & Mendell (2000) computed ordinary fluid-like r -modes, taking account of the entrainment effects in the core. They showed that the mutual friction damping in the core is not effective to suppress the instability except for a few domains of the entrainment parameter η . Recently, Lee & Yoshida (2002) have studied the r -modes in neutron stars with the superfluid core with a different method, confirming most of the results by Lindblom & Mendell (2000). Lee & Yoshida (2002) have also found that, as suggested by Andersson & Comer (2001), the r -modes are split into ordinary fluid-like r -modes and superfluid-like r -modes, and that the instability caused by the superfluid-like r -modes is extremely weak and easily damped by dissipation processes in the interior.

The purpose of this paper is to study modal properties of inertial modes in rotating neutron stars containing superfluids in the core. As shown by Lockitch & Friedman (1999), the existence of inertial modes in the lowest order of angular frequency Ω of rotation depends on the structure of the so-called zero-frequency subspace of eigensolutions to linearized hydrodynamic equations. Naively speaking, we may say that, for inertial modes to exist in the limit of $\Omega \rightarrow 0$, g -modes of a non-rotating star with $\Omega = 0$ need to be degenerate to the zero-frequency subspace. Investigating the zero-frequency subspace of pulsation equations for the superfluid core, Andersson & Comer (2001) showed the existence of the zero-frequency g -modes, and hence the existence of inertial modes (see, also, Comer 2002). In this paper, we calculate global inertial mode oscillations in neutron stars with a core filled with neutron and proton superfluids, taking account of the entrainment effects between the two superfluids. For the dynamics of superfluids in the core, we employ a two-fluid formalism similar to that given in Lindblom & Mendell (1994). To obtain the inertial modes in the limit of $\Omega \rightarrow 0$, we generalize the method devised by Lockitch & Friedman (1999) to stars composed of a superfluid core and a normal fluid envelope. In §2 we present the basic equations employed in this paper for the dynamics of superfluids in the core, and in §3 dissipation processes considered in this paper are described. §4 gives numerical results, and §5 and §6 are for discussions and conclusions.

2 OSCILLATION EQUATIONS

We derive basic equations governing superfluid motions in the neutron star core in the Newtonian dynamics, assuming uniform rotation of the star. The core is assumed to be filled with neutron and proton superfluids and a normal fluid of electron. We also assume perfect charge neutrality between the protons and electrons because the plasma frequency is much higher than the oscillation frequencies considered in this paper (see, e.g., Mendell 1991a). Since the transition temperatures $T_c \sim 10^9$ K to neutron and proton superfluids are much higher than the interior temperatures of old neutron stars (see, e.g., Epstein 1988), we assume that all the neutrons and protons in the core are in superfluid states and the excited normal fluid components of the fluids can be ignored.

In this study, we assume that the neutron and proton superfluids and the electron normal fluid in the equilibrium state are in the same rotational motion with the angular velocity Ω around the axis of rotation. The velocity of the fluid v^a in the equilibrium state is therefore given by $v^a = \Omega \varphi^a$, where φ^a is the rotational Killing vector. In a perturbed state, however, the

neutron and the proton superfluids can move differently from each other in the core, obeying their own governing equations. In order to describe the perturbations in the superfluid core, thus, we can choose the perturbed neutron density $\delta\rho_n$, the perturbed proton density $\delta\rho_p$, the perturbed velocity of the neutron δv_n^a , and the perturbed velocity of the proton δv_p^a as dynamical variables. Since the equilibrium star is axisymmetric about the rotation axis, we assume that the time and azimuthal dependence of the perturbations is given by $\exp(i\sigma t + im\varphi)$ with σ being the oscillation frequency observed in an inertial frame, and m is an integer representing the azimuthal wave-number. The basic perturbation equations employed in this paper for the neutron and proton superfluids in rotating neutron stars are essentially the same as those given in Mendell (1991a) and Lindblom & Mendell (1994) and are given as follows:

$$i\omega\delta\rho_n + \nabla_a(\rho_n\delta V_n^a) = 0, \quad (1)$$

$$i\omega\delta\rho_p + \nabla_a(\rho_p\delta V_p^a) = 0, \quad (2)$$

$$i\omega\delta v_n^a + 2\Omega\delta v_n^b\nabla_b\varphi^a + \nabla^a(\delta\mu_n + \delta\Phi) - 2\Omega\frac{\rho_{np}}{\rho_n}(\delta v_p^b - \delta v_n^b)\nabla^a\varphi_b = 0, \quad (3)$$

$$\begin{aligned} & i\omega\left(\delta v_p^a + \frac{m_e}{m_p}\delta v_e^a\right) + 2\Omega\left(\delta v_p^b + \frac{m_e}{m_p}\delta v_e^b\right)\nabla_b\varphi^a \\ & + \nabla^a\left(\delta\mu_p + \frac{m_e}{m_p}\delta\mu_e + \left(1 + \frac{m_e}{m_p}\right)\delta\Phi\right) - 2\Omega\frac{\rho_{np}}{\rho_n}(\delta v_n^b - \delta v_p^b)\nabla^a\varphi_b = 0, \end{aligned} \quad (4)$$

where ω denotes the corotating frequency, defined by $\omega = \sigma + m\Omega$, ∇_a means the covariant derivative in flat three dimensional space, and δQ stands for the Eulerian change in the physical quantity Q . Here, m_p and m_e are the proton and the electron masses. In the mass conservation equations (1) and (2), $\rho_n\delta V_n^a$ and $\rho_p\delta V_p^a$ mean the mass current vector of the neutron and of the proton, respectively, and can be expanded in terms of the perturbations of the superfluid velocities as follows:

$$\rho_n\delta V_n^a = \rho_{nn}\delta v_n^a + \rho_{np}\delta v_p^a, \quad (5)$$

$$\rho_p\delta V_p^a = \rho_{pn}\delta v_n^a + \rho_{pp}\delta v_p^a, \quad (6)$$

where the relationships $\rho_n = \rho_{nn} + \rho_{np}$, $\rho_p = \rho_{pn} + \rho_{pp}$, and $\rho_{np} = \rho_{pn}$ must be satisfied due to the Galilean invariance of equations (1) and (2). The perturbed superfluid velocities are then given in terms of δV_n^a and δV_p^a as

$$\delta v_n^a = \frac{\rho_{11}}{\rho_n}\delta V_n^a + \frac{\rho_{12}}{\rho_n}\delta V_p^a, \quad (7)$$

$$\delta v_p^a = \frac{\rho_{21}}{\rho_p}\delta V_n^a + \frac{\rho_{22}}{\rho_p}\delta V_p^a, \quad (8)$$

where

$$\rho_{11} = \frac{\rho_{pp}\rho_n^2}{\tilde{\rho}^2}, \quad \rho_{22} = \frac{\rho_{nn}\rho_p^2}{\tilde{\rho}^2}, \quad \rho_{12} = \rho_{21} = -\frac{\rho_{np}\rho_n\rho_p}{\tilde{\rho}^2}, \quad (9)$$

and

$$\tilde{\rho}^2 = \rho_{nn}\rho_{pp} - \rho_{np}\rho_{pn}. \quad (10)$$

Note that $\rho_{11} + \rho_{12} = \rho_n$ and $\rho_{22} + \rho_{21} = \rho_p$. In Euler equations (3) and (4), δv_e^a and $\delta\Phi$ are the perturbed velocity of the electron and the perturbed gravitational potential, respectively, and $\delta\mu_n$, $\delta\mu_p$, and $\delta\mu_e$ stand for the perturbed chemical potentials per unit mass for the neutron, proton and electron, respectively. Because of the assumption of the perfect charge neutrality in the proton and electron plasma, we may have

$$\delta V_p^a = \delta v_e^a, \quad \text{and} \quad \delta\rho_p/m_p = \delta\rho_e/m_e, \quad (11)$$

where ρ_e is the electron density. The perturbed gravitational potential is determined by the Poisson equation

$$\nabla^2\delta\Phi = 4\pi G\delta\rho, \quad (12)$$

where $\rho = \rho_n + \rho_p + \rho_e$ and G is the gravitational constant.

Using a variant of Gibbs-Duhem relation, the pressure perturbation is given by

$$\delta p = \rho_n\delta\mu_n + \rho_p\delta\tilde{\mu}, \quad (13)$$

where $\delta\tilde{\mu} = \delta\mu_p + \frac{m_e}{m_p}\delta\mu_e$. Here, we have ignored the entropy carried by the electron normal fluid. This treatment is justified for a star whose temperature is low enough. Note also that the superfluids carry no entropy.

To obtain a relation between the densities ρ_n and ρ_p and the chemical potentials μ_n and $\tilde{\mu}$, we begin with writing the energy density e as

$$e = e(\rho_n, \rho_p), \quad (14)$$

with which the chemical potentials are defined as

$$\mu_n(\rho_n, \rho_p) = (\partial e / \partial \rho_n)_{\rho_p}, \quad \mu_p(\rho_n, \rho_p) = (\partial e / \partial \rho_p)_{\rho_n}. \quad (15)$$

If the chemical potential of the electron is given by

$$\mu_e(\rho_e) = c^2 \sqrt{1 + (3\pi^2 \hbar^3 \rho_e / m_e^4 c^3)^{2/3}}, \quad (16)$$

we have, due to charge neutrality condition $\rho_e = \rho_p m_e / m_p$,

$$\begin{pmatrix} \delta \mu_n \\ \delta \tilde{\mu} \end{pmatrix} = \begin{pmatrix} \mathcal{P}_{11} & \mathcal{P}_{12} \\ \mathcal{P}_{21} & \mathcal{P}_{22} \end{pmatrix} \begin{pmatrix} \delta \rho_n \\ \delta \rho_p \end{pmatrix}, \quad (17)$$

where

$$\mathcal{P}_{11} = \left(\frac{\partial \mu_n}{\partial \rho_n} \right)_{\rho_p}, \quad \mathcal{P}_{12} = \left(\frac{\partial \mu_n}{\partial \rho_p} \right)_{\rho_n} = \left(\frac{\partial \tilde{\mu}}{\partial \rho_n} \right)_{\rho_p} = \mathcal{P}_{21}, \quad \mathcal{P}_{22} = \left(\frac{\partial \tilde{\mu}}{\partial \rho_p} \right)_{\rho_n}. \quad (18)$$

We write the inverse of equation (17) as

$$\begin{pmatrix} \delta \rho_n \\ \delta \rho_p \end{pmatrix} = \begin{pmatrix} \mathcal{Q}_{11} & \mathcal{Q}_{12} \\ \mathcal{Q}_{21} & \mathcal{Q}_{22} \end{pmatrix} \begin{pmatrix} \delta \mu_n \\ \delta \tilde{\mu} \end{pmatrix}, \quad (19)$$

where

$$\begin{pmatrix} \mathcal{Q}_{11} & \mathcal{Q}_{12} \\ \mathcal{Q}_{21} & \mathcal{Q}_{22} \end{pmatrix} = \begin{pmatrix} \mathcal{P}_{11} & \mathcal{P}_{12} \\ \mathcal{P}_{21} & \mathcal{P}_{22} \end{pmatrix}^{-1}. \quad (20)$$

The curl of the Euler equations, which is the vorticity equation, is useful for perturbation analysis of rotating stars, and is given in terms of δV_n^a and δV_p^a by

$$\epsilon^{abc} \nabla_b \left[\left(i\omega \frac{\rho_{11}}{\rho_n} g_{cd} + 2\Omega \nabla_d \varphi_c \right) \delta V_n^d + i\omega \frac{\rho_{12}}{\rho_n} g_{cd} \delta V_p^d \right] = 0, \quad (21)$$

$$\epsilon^{abc} \nabla_b \left[\left(i\omega \left(\frac{\rho_{22}}{\rho_p} + \frac{m_e}{m_p} \right) g_{cd} + 2\Omega \left(1 + \frac{m_e}{m_p} \right) \nabla_d \varphi_c \right) \delta V_p^d + i\omega \frac{\rho_{12}}{\rho_p} g_{cd} \delta V_n^d \right] = 0, \quad (22)$$

where g_{ab} and ϵ^{abc} mean the metric tensor and the Levi-Civita tensor in the flat space.

Although we have made no assumptions for rotation velocity up to now, let us restrict our consideration to slowly rotating stars in order to obtain inertial modes to the lowest order in the stellar rotation frequency Ω . In the limit of slow rotation, then, we assume perturbations to obey the following ordering in Ω :

$$\delta V_n^a = O(1), \quad \delta V_p^a = O(1), \quad \omega/\Omega = O(1), \quad \delta \rho_n = O(\Omega), \quad \delta \rho_p = O(\Omega), \quad \text{and} \quad \delta \Phi = O(\Omega). \quad (23)$$

Examining the zero-frequency subspace of the perturbation in a non-rotating superfluid core, Andersson & Comer (2001) showed that solutions obeying the ordering (23) may be allowed in the equations describing hydrodynamics in the superfluid core. The existence of solutions obeying (23) is similar to that in a barotropic normal fluid star (Lockitch & Friedman 1999). Those solutions can be interpreted as infinitely degenerate g -modes of zero-frequency if we consider no rotational effect. It is convenient to introduce dimensionless frequency κ_0 in the limit of $\Omega \rightarrow 0$, which is defined by

$$\kappa_0 = \lim_{\Omega \rightarrow 0} \omega/\Omega. \quad (24)$$

Then, basic equations describing the lowest order inertial modes are given by

$$\nabla_a (\rho_n \delta V_n^a) = 0, \quad \nabla_a (\rho_p \delta V_p^a) = 0, \quad (25)$$

$$q_n^a = \epsilon^{abc} \nabla_b \left[\left(\frac{\rho_{11}}{\rho_n} g_{cd} - 2i\kappa_0^{-1} \nabla_d \varphi_c \right) \delta V_n^d + \frac{\rho_{12}}{\rho_n} g_{cd} \delta V_p^d \right] = 0, \quad (26)$$

$$q_p^a = \epsilon^{abc} \nabla_b \left[\left(\left(\frac{\rho_{22}}{\rho_p} + \frac{m_e}{m_p} \right) g_{cd} - 2i\kappa_0^{-1} \left(1 + \frac{m_e}{m_p} \right) \nabla_d \varphi_c \right) \delta V_p^d + \frac{\rho_{12}}{\rho_p} g_{cd} \delta V_n^d \right] = 0. \quad (27)$$

Note that q_n^a (q_p^a) has only two independent components because $\nabla_a q_n^a = 0$ ($\nabla_a q_p^a = 0$) is automatically satisfied.

From similar consideration to that for the superfluid core, basic equations for pulsations in the normal fluid envelope are given by

$$\nabla_a (\rho \delta v^a) = 0, \quad (28)$$

$$q^a = \epsilon^{abc} \nabla_b [(g_{cd} - 2i\kappa_0^{-1} \nabla_d \varphi_c) \delta v^d] = 0, \quad (29)$$

where δv^a denotes the perturbation of the fluid velocity.

The perturbations in a uniformly rotating star are expanded in terms of spherical harmonic functions with different l 's for a given m (e.g., Lee & Saio 1986). The vector δV_X^a is therefore given by

$$\delta V_X^r = r \sum_{l \geq |m|} S_l^X(r) Y_l^m(\theta, \varphi) e^{i\sigma t}, \quad (30)$$

$$\delta V_X^\theta = \sum_{l \geq |m|} \left[H_l^X(r) \frac{\partial}{\partial \theta} Y_l^m(\theta, \varphi) - iT_{l'}^X(r) \frac{1}{\sin \theta} \frac{\partial}{\partial \varphi} Y_{l'}^m(\theta, \varphi) \right] e^{i\sigma t}, \quad (31)$$

$$\delta V_X^\varphi = \frac{1}{\sin^2 \theta} \sum_{l \geq |m|} \left[H_l^X(r) \frac{\partial}{\partial \varphi} Y_l^m(\theta, \varphi) + iT_{l'}^X(r) \sin \theta \frac{\partial}{\partial \theta} Y_{l'}^m(\theta, \varphi) \right] e^{i\sigma t}, \quad (32)$$

where $l = |m| + 2(k-1)$ and $l' = l + 1$ for even modes, and $l = |m| + 2k - 1$ and $l' = l - 1$ for odd modes, where $k = 1, 2, 3, \dots$. Here, we have employed the constituent index X which can be either n , p or o , where o means the velocity field in the normal fluid envelope, and the normal spherical polar coordinate (r, θ, φ) is used. In this paper, the even and odd modes are respectively characterized by symmetry and antisymmetry of their eigenfunctions with respect to the equatorial plan. Substituting these expansions into equations (25) through (27), we obtain oscillation equations given as a set of simultaneous linear ordinary differential equations of the expansion coefficients (see Appendix), which is to be integrated in the superfluid core. The oscillation equations solved in the normal fluid envelope are similar to those given in Lockitch & Friedman (1999).

To obtain a unique solution of an oscillation mode, solutions in the superfluid core and in the normal fluid envelope are matched at the interface between the two domains by imposing jump conditions given by

$$\delta V_n^r = \delta V_p^r = \delta V_o^r. \quad (33)$$

The boundary conditions at the stellar center is the regularity condition of the perturbations δV_n^a and δV_p^a . The boundary condition at the stellar surface is $\Delta p = 0$, where Δp is the Lagrangian change in the pressure.

For numerical computation, oscillation equations of a finite dimension are obtained by disregarding the terms with l larger than $l_{max} = |m| + 2k_{max} - 1$ in the expansions of perturbations such as given by (30) to (32), where l_{max} is determined so that the eigenfrequency and the eigenfunctions are well converged as l_{max} increases. We solve the oscillation equations of a finite dimension as an eigenvalue problem with the scaled oscillation frequency κ_0 using a Henyey type relaxation method (see, e.g., Unno et al. 1989).

3 DISSIPATIONS

In the previous section, we consider no dissipation effects. Thus, the conserved energy E of oscillations exists and is given by (Mendell 1991b)

$$\begin{aligned} E = & \frac{1}{2} \int_{V_S} \left(\rho \delta v_a^* \delta v^a + \frac{\tilde{\rho}^2}{\rho} \delta w_a^* \delta w^a + \sum_{i,j} \mathcal{P}_{ij} \delta \rho_i^* \delta \rho_j - \frac{1}{4\pi G} \nabla_a \delta \Phi^* \nabla^a \delta \Phi \right) \sqrt{g} d^3 x \\ & + \frac{1}{2} \int_{V_N} \left(\rho \delta v_a^* \delta v^a + \left(\frac{\delta p}{\rho} + \delta \Phi \right) \delta \rho^* \right) \sqrt{g} d^3 x \end{aligned} \quad (34)$$

where g is the determinant of the metric, and asterisk (*) denotes the complex conjugate of the quantity. Here, $\int_{V_S} d^3 x$ and $\int_{V_N} d^3 x$ means the integration over the superfluid core and the normal fluid envelope, respectively. In this equation, we have employed new variables, defined by

$$\begin{aligned} \delta v^a &= (\rho_n \delta v_n^a + \rho_p \delta v_p^a) / \rho = (\rho_n \delta V_n^a + \rho_p \delta V_p^a) / \rho, \\ \delta w^a &= \delta v_n^a - \delta v_p^a = \frac{\rho_n \rho_p}{\tilde{\rho}^2} (\delta V_n^a - \delta V_p^a), \\ \delta \rho_1 &= \delta \rho_n, \quad \delta \rho_2 = \delta \rho_p. \end{aligned} \quad (35)$$

Note that $\rho \delta v^a$ is interpreted as the total momentum current vector of perturbation. In reality, however, there are a lot of dissipation mechanisms in real neutron stars. Thus, the oscillation energy E is not conserved in real neutron stars. In other words, the amplitude of an oscillation mode should damp or grow depending on considered dissipation mechanisms.

The stability of an oscillation mode of a star is then determined by summing up all contributions from various dissipation mechanisms. In this study, we consider the contributions from gravitational radiation reaction, viscous processes, and mutual friction in the superfluid core. Then, the averaged energy loss (or gain) rate dE/dt of a normal mode in a rotating neutron star may be given by

$$\begin{aligned} \frac{dE}{dt} &= -\sigma\omega \sum_{l' \geq |m|} N_l \sigma^{2l} |\delta J_{l'm}|^2 - 2 \int_{V_S+V_N} \eta_s \delta \sigma^{ab*} \delta \sigma_{ab} \sqrt{g} d^3x \\ &\quad - 2\Omega \int_{V_S} \rho_n B_n \left(\frac{\tilde{\rho}^2}{\rho_n \rho_p} \right)^2 (g_{ab} - z_a z_b) \delta w^{a*} \delta w^b \sqrt{g} d^3x \\ &= \sum_{l' \geq |m|} \left(\frac{dE}{dt} \right)_{GJ,l'} + \left(\frac{dE}{dt} \right)_S + \left(\frac{dE}{dt} \right)_{MF}, \end{aligned} \quad (36)$$

where z^a is the unit vector that parallels the rotation axis, defined by $z_a = \epsilon_{abc} \nabla^b \varphi^c / 2$.

The terms $(dE/dt)_{GJ,l}$ on the right hand side of equation (36) denote the energy loss (or gain) rates due to gravitational radiation reaction associated with the mass current multipole moment δJ_{lm} , where

$$\delta J_{lm} = \frac{2}{c(l+1)} \int_{V_S+V_N} r^{l+1} \rho \delta v_a \epsilon^{abc} \nabla_b Y_l^{m*} \nabla_c r \sqrt{g} d^3x \quad (37)$$

and

$$N_l = \frac{4\pi G}{c^{2l+1}} \frac{(l+1)(l+2)}{l(l-1)[(2l+1)!!]^2}, \quad (38)$$

and c is the velocity of light (Thorne 1980, Lindblom et al. 1998).

The term $(dE/dt)_S$ is the energy dissipation rate due to shear viscosity, and η_s is the shear viscosity coefficient, and $\delta \sigma_{ab}$ is the traceless rate of strain tensor for the perturbed velocity field, which is defined by

$$\delta \sigma_{ab} = \frac{1}{2} \left(\nabla_a u_b + \nabla_b u_a - \frac{2}{3} g_{ab} \nabla_c u^c \right), \quad (39)$$

for the velocity field u^a . In this paper, we ignore the contribution from the bulk viscosity, which is important only for newly born hot neutron stars without superfluids in the core. The shear viscosity coefficient we use in the superfluid core is

$$\eta_s = 6.0 \times 10^{18} \left(\frac{\rho}{10^{15} \text{g/cm}^3} \right)^2 \left(\frac{10^9 \text{K}}{T} \right)^2 \text{g/cm s} \quad (40)$$

(Cutler & Lindblom 1987, Flowers & Itoh 1979), and that in the normal fluid envelope is given by

$$\eta_S = 1.95 \times 10^{18} \left(\frac{\rho}{10^{15} \text{g/cm}^3} \right)^{9/4} \left(\frac{10^9 \text{K}}{T} \right)^2 \text{g/cm s} \quad (41)$$

(Cutler & Lindblom 1987, Flowers & Itoh 1979). The rate of strain $\delta \sigma_{ab}$ is evaluated by using δv_e^a in the superfluid core, and δv^a in the normal fluid envelope (Lindblom & Mendell 2000).

The term $(dE/dt)_{MF}$ is the energy loss rate due to the mutual friction in the superfluid core, and the dimensionless coefficient B_n is given by (Mendell 1991b)

$$B_n = 0.011 \times \frac{\rho_p}{\rho_n} \left(\frac{\rho_{pp}}{\rho_p} \right)^{1/2} \left(\frac{\rho_{pn}}{\rho_{pp}} \right)^2 \left(\frac{\rho_p}{10^{14} \text{g/cm}^3} \right)^{1/6}. \quad (42)$$

The mutual friction is a dissipation mechanism inherent to a rotating system of superfluids, and it is caused by scattering of normal fluid particles off the vortices in the superfluids. Since we have assumed perfect charge neutrality between the electrons and protons, we consider scattering between the normal electrons and vortices of the neutron superfluid (Mendell 1991b, see also Alpar, Langer, & Sauls, 1984).

As is indicated by the first term on the right hand side of equation (36), if the normal mode has an oscillation frequency that satisfies $-\sigma\omega > 0$, the oscillation energy E in the corotating frame, in the absence of other damping mechanisms, increases as a result of gravitational wave radiation, indicating instability of the mode (Friedman & Schutz 1978).

The damping (or growth) time-scale τ of a normal mode may be given by

$$\frac{1}{\tau} = -\frac{1}{2E} \left(\frac{dE}{dt} \right) = \sum_{l' > |m|} \frac{1}{\tau_{GJ,l'}} + \frac{1}{\tau_S} + \frac{1}{\tau_{MF}}, \quad (43)$$

where $\tau_i = -2E/(dE/dt)_i$. For the inertial modes, it is convenient to derive an extrapolation formula of the time-scale τ given as a function of Ω and the interior temperature T (e.g., Lindblom et al. 1998, Lindblom & Mendell 2000):

$$\frac{1}{\tau(\Omega, T)} = \sum_{l' \geq |m|} \frac{1}{\tau_{GJ, l'}^0} \left(\frac{\Omega^2}{\pi G \bar{\rho}} \right)^{l'} + \frac{1}{\tau_S^0} \left(\frac{10^9 K}{T} \right)^2 + \frac{1}{\tau_{MF}^0} \left(\frac{\Omega^2}{\pi G \bar{\rho}} \right)^{1/2}, \quad (44)$$

where $\bar{\rho} = M/(4\pi R^3/3)$. Here, M and R are the mass and radius of the star. The quantities $\tau_{GJ, l'}^0$, τ_S^0 , and τ_{MF}^0 are assumed to be independent on Ω and T .

4 NUMERICAL RESULTS

Following Lindblom & Mendell (2000), we employ a polytropic model of index $N = 1$ with the mass $M = 1.4M_\odot$ and the radius $R = 12.57\text{km}$ as a background model for modal analysis. The model is divided into a superfluid core and a normal fluid envelope, the interface of which is set at $\rho = \rho_s = 2.8 \times 10^{14} \text{g/cm}^3$. In the normal fluid envelope, we use the polytropic equation of state given by $p = K\rho^2$ both for the equilibrium structure and for the oscillation equations. The core is assumed to be filled with neutron and proton superfluids and a normal fluid of electron, for which an equation of state, labeled A18+ δv +UIX (Akmal, Pandharipande, & Ravenhall 1998), is used to give the thermodynamical quantities used for the oscillation equations. For the mass density coefficients ρ_{nn} , ρ_{pp} , and ρ_{np} in the core, we employ an empirical relation given by (see Lindblom & Mendell 2000)

$$\rho_p/\rho \approx 0.031 + 8.8 \times 10^{-17} \rho, \quad (45)$$

and a formula given by

$$\rho_{np} = -\eta \rho_n, \quad (46)$$

where η is a parameter of order of ~ 0.04 (Borumand, Joynt, & Kluźniak 1996). In this paper we call η the entrainment parameter. In this study, only the modes having $m = 2$ are examined because they are the most significant for the gravitational radiation driven instability. For the range of the entrainment parameter η , we choose $-0.01 \leq \eta \leq 0.04$. Although negative values of η might not be suitable for physical models of neutron stars, we include in our calculations negative η to demonstrate clearly the η dependence of the inertial modes around $\eta \sim 0$.

We find that inertial modes in the superfluid core are split into two families, which we call ordinary fluid inertial modes (i^o -mode), and superfluid inertial modes (i^s -mode). This mode splitting is common in other oscillation modes in the superfluid core, because the dynamical degree of freedom in the superfluid core simply doubles (see, e.g., Epstein 1988; Lindblom & Mendell 1994; Lee 1995; Andersson & Comer 2001; Prix & Rieutord 2002). As a subclass of inertial modes, the r -modes of neutron stars with a superfluid core are also split into ordinary fluid-like r -modes and superfluid-like r -modes, which we respectively call r^o -modes and r^s -modes as in Lee & Yoshida (2002). Since basic properties of the eigenfunctions δv^a (or δw^a) of an inertial mode are quite similar to the eigenfunctions δv^a of the corresponding inertial mode in the normal fluid star without core superfluidity, we employ the same classification scheme as that employed by Lockitch & Friedman (1999) and Yoshida & Lee (2000a) for inertial modes in the normal fluid star. In this scheme, inertial modes are classified in terms of the two quantum numbers m and $l_0 - |m|$, and the scaled frequency $\kappa_0 \equiv \lim_{\Omega \rightarrow 0} \omega/\Omega$. For given m and $l_0 - |m|$, the number of inertial modes of one kind is $l_0 - |m|$. Even parity inertial modes have even $l_0 - |m|$, and odd parity inertial modes have odd $l_0 - |m|$. Note that the r -modes are classified as inertial modes with $l_0 - |m| = 1$ (e.g., Yoshida & Lee 2000a). It is interesting to note that, although there exist three quantum numbers to classify the normal modes of non-rotating stars, a third quantum number may be missing for inertial modes. Usually, the ratio E_S/E of the oscillation energy E_S in the core to the total oscillation energy E is a good indicator for a distinction between i^s - and i^o -modes, since we find $E_S/E \lesssim 0.5$ for the i^o -modes and $E_S/E \gtrsim 0.9$ for the i^s -modes for the models composed of a superfluid core and a normal fluid envelope.

Let us first consider the case of $\eta = 0$, that is, the case of no entrainment. In Table 1, κ_0 's for i^o - and i^s -modes of low $l_0 - |m|$ are tabulated together with κ_0 's for the corresponding inertial modes in the $N = 1$ polytrope model without superfluidity in the core, which we simply call i -mode (see, Lockitch & Friedman 1999; Yoshida & Lee 2000a). Our values of κ_0 for the i^o - and i^s -modes are in good agreement with those computed by Lee & Yoshida (2002) with a different method. Note that we have shown five digits for numerical values of κ_0 to indicate that our numerical results include at most a few percent numerical error. We can see that κ_0 's of paired i^o - and i^s -modes have quite similar values to that of the corresponding inertial mode in the $N = 1$ polytropic normal fluid model. In order to demonstrate model properties of the inertial modes, we exhibit the expansion coefficients S_l^v , $T_{l'}^v$, S_l^w , and $T_{l'}^w$ of the eigenfunctions δv^a and δw^a , defined by

$$S_l^v = (\rho_n S_l^n + \rho_p S_l^p)/\rho, \quad T_{l'}^v = (\rho_n T_{l'}^n + \rho_p T_{l'}^p)/\rho, \quad (47)$$

$$S_l^w = \rho_n \rho_p (S_l^n - S_l^p)/\bar{\rho}^2, \quad T_{l'}^w = \rho_n \rho_p (T_{l'}^n - T_{l'}^p)/\bar{\rho}^2. \quad (48)$$

Note that S_l^w and $T_{l'}^w$ vanish in the normal fluid envelope because neutrons and protons co-move. In Figures 1 through 4, the first three expansion coefficients S_l^v , $T_{l'}^v$, S_l^w , and $T_{l'}^w$ are shown for the i^o - and i^s -modes of $l_0 - |m| = 2$ which have $\kappa_0 = -0.5685$ and $\kappa_0 = -0.5115$, respectively. Here, the amplitude normalization given by $T_{m+1}^v = 1$ has been used. It is

noted that δv^a of the i^o -modes and δw^a of the i^s -modes resemble δv^a of the corresponding inertial modes of the normal fluid star model (compare the figures with Figure 1 of Yoshida & Lee 2000a). It is also found that although the amplitudes of δv^a and δw^a are similar to each other for the i^o -modes, the amplitude of δw^a dominates that of δv^a for the i^s -modes, which indicates that the two superfluids in the core basically counter-move for the i^s -modes.

In Figures 5 to 8, κ_0 's of inertial modes of low $l_0 - |m|$ are plotted as functions of the entrainment parameter η , where Figure 5 is for even parity modes with positive κ_0 , Figure 6 for even parity modes with negative κ_0 , Figure 7 for odd parity modes with positive κ_0 , and Figure 8 for odd parity modes with negative κ_0 . In Figure 7, we have also given κ_0 of the r^s -mode, the locus of which as a function of η is in good agreement with the locus calculated by Lee & Yoshida (2002) with a different method. Note that we cannot obtain r^o -modes by using our numerical method, the reason for which is not clear (see the next section)¹. From Figures 5 through 8, we find that, although κ_0 for the i^o -modes is only weakly dependent on the parameter η , $|\kappa_0|$ for the i^s -modes increases almost linearly as η increases. Because of these different dependences of κ_0 on the parameter η , frequent avoided crossings are inevitable between i^o -modes and i^s -modes. As shown by Figure 7, for example, the i^s -mode of $l_0 - |m| = 3$ which has $\kappa_0 = 1.3865$ at $\eta = 0$ suffers three avoided crossings with different i^o -modes as η is increased from -0.01 to 0.04 . Since κ_0 's of paired i^o - and i^s -modes at $\eta = 0$ are close to each other, they always make an avoided crossing around $\eta = 0$, which is also shown by Figures 5 through 8.

In Figures 9 through 12, we display the first three coefficients S_l^v , T_l^v , S_l^w , and T_l^w for the i^o - and i^s -modes of $l_0 - |m| = 2$, which respectively have $\kappa_0 = -0.5403$ and $\kappa_0 = -0.6103$ at $\eta = 0.01$. Here, the amplitude normalization given by $T_{m+1}^v = 1$ has been used. Note that these two inertial modes have $\kappa_0 = -0.5685$ and $\kappa_0 = -0.5115$ at $\eta = 0$. Comparing these figures with Figures 1 to 4, we observe that the basic properties of the eigenfunctions of the inertial modes do not depend on the entrainment parameter η , except at avoided crossings.

In Table 2, we tabulate κ_0 and the scaled damping (or growth) timescales τ_i^0 for the i^o - and i^s -modes with positive κ_0 for $\eta = 0, 0.02$, and 0.04 . We find that the growth timescales $\tau_{GJ,l}^0$ for all the i^o - and i^s -modes in the table are longer than that of the r -mode of the normal fluid neutron star, which means that the instability of the i^o - and i^s -modes caused by the current multipole radiation is not strong. This result is the same as that for the inertial modes in a normal fluid star. It is interesting to note that the two i^s -modes in the table having $\kappa_0 = 2.1661$ and $\kappa_0 = 2.4083$ at $\eta = 0.04$ are stable against gravitational radiation reaction since they have positive $\tau_{GJ,l}^0$. The reason for this can be understood as follows. As mentioned in the last section, the gravitational radiation driven instability sets in when the frequency of the mode satisfies $\sigma(\sigma - m\Omega) < 0$, which leads to the instability criterion given by $0 < \kappa_0 < m$ for positive m . For inertial modes of normal fluid stars, it is known that all inertial modes having positive κ_0 for $m > 0$ satisfy the instability criterion (Lindblom & Ipser 1999; Lockitch & Friedman 1999; Yoshida & Lee 2000a). For inertial modes in the superfluid core, on the other hand, since the frequency $|\kappa_0|$ of the i^s -modes monotonically increases as the value of η is increased, the κ_0 of some positive-frequency i^s -modes exceeds the limit m when η gets larger than a critical value η_{crit} , the value of which depends on the mode (see, Figures 5 and 7). Thus, some of the positive-frequency i^s -modes that are unstable satisfying the instability criterion at $\eta = 0$ become stable invalidating the criterion for η larger than η_{crit} . As for the effects of mutual friction, we observe that the damping timescales τ_{MF}^0 for the inertial modes are much shorter than those due to other dissipation processes unless mutual friction between the superfluids is extremely weak, that is, the parameter η is extremely small. Note that the amount of dissipation caused by the mutual friction is roughly proportional to $B_n |\delta w^a|^2$ where $B_n \propto \eta^2$. Accordingly, if we assume $\eta \sim 0.04$ as a typical value for the parameter, the dissipation due to mutual friction in the superfluid core is very effective to damp out the i^o - and i^s -modes.

5 DISCUSSION

As mentioned in the previous section, with our method of calculation we fail in obtaining ordinary fluid r -modes, which are called r^o -modes in Lee & Yoshida (2002). However, this does not necessarily mean that our basic equations (A2)–(A7) do not contain such r -mode solutions. To prove this, let us consider purely axial parity solutions, whose angular behavior is given by a purely axial harmonic function having $l' = m$. In this section, we also assume for simplicity that the densities ρ_n , ρ_p , and ρ_{12} are constant and do not depend on r . In virtue of the assumption of purely axial mode, we can reduce our basic equations to the following equations (see, Appendix):

For the superfluid core,

¹ Recently, we have generalized our formalism to the case of general relativistic superfluid neutron stars. In that study, our numerical code for Newtonian dynamics have been straightforwardly extended into the one for general relativistic dynamics. With this relativistic version of the code, both r^o - and r^s -modes can be obtained even in almost the Newtonian regime with small values of the relativistic parameter M/R . Fundamental differences in the two formalisms reside only in the master equations. As shown in equations (49)–(52), the basic equations for neutron and proton fluids in the Newtonian case get to degenerate in lowest order of Ω when r^o -modes are considered, while the basic equations in the relativistic case don't even if we are concerned with r^o -modes (see, Yoshida & Lee 2003). We therefore guess that the reason why r^o -modes cannot be obtained in the Newtonian case is this possible degeneracy of the two sets of superfluid equations in Newtonian dynamics.

$$\nu m(m+2)J_{m+1}^m \left\{ r \frac{d}{dr} T_m^n - (m-1)T_m^n \right\} = 0, \quad (49)$$

$$\nu m(m+2)J_{m+1}^m \left\{ r \frac{d}{dr} T_m^p - (m-1)T_m^p \right\} = 0, \quad (50)$$

$$\left(1 - \frac{m\nu}{m(m+1)} \right) T_m^n + \frac{\rho_{12}}{\rho_n} (T_m^p - T_m^n) = 0, \quad (51)$$

$$\left(1 - \frac{m\nu}{m(m+1)} \right) T_m^p + \frac{\rho_{12}}{\rho_p} (T_m^n - T_m^p) = 0, \quad (52)$$

For the normal fluid envelope,

$$\nu m(m+2)J_{m+1}^m \left\{ r \frac{d}{dr} T_m^o - (m-1)T_m^o \right\} = 0, \quad (53)$$

$$\left(1 - \frac{m\nu}{m(m+1)} \right) T_m^o = 0, \quad (54)$$

where $\nu = 2\Omega/\sigma$ and $J_l^m = [(l^2 - m^2)/(4l^2 - 1)]^{1/2}$. Here, we have ignored the terms due to m_e/m_p for simplicity. The condition for the continuity of the total momentum flux at the interface between the superfluid core and the normal fluid envelope is given by

$$T_m^o = (\rho_n T_m^n + \rho_p T_m^p)/\rho \quad \text{at } r = r_s, \quad (55)$$

where r_s is the radius of the superfluid core. Because the factor $\nu m(m+2)J_{m+1}^m$ in equations (49), (50) and (53) does not vanish for $\nu \neq 0$ and $m \geq 2$, we can obtain solutions to differential equations (49), (50), and (53) as

$$T_m^n = A_n(r/r_s)^{m-1}, \quad T_m^p = A_p(r/r_s)^{m-1}, \quad T_m^o = B(r/r_s)^{m-1}, \quad (56)$$

where A_n , A_p , and B are constants, which are to be determined by boundary condition (55) and constraint equations (51), (52), (54). Substituting the solutions (56) into algebraic equations (51), (52), (54) and (55), we obtain

$$\begin{pmatrix} f - \rho_{12}/\rho_n & \rho_{12}/\rho_n \\ \rho_{12}/\rho_p & f - \rho_{12}/\rho_p \end{pmatrix} \begin{pmatrix} A_n \\ A_p \end{pmatrix} \equiv W \begin{pmatrix} A_n \\ A_p \end{pmatrix} = 0, \quad (57)$$

$$fB = 0, \quad (58)$$

and

$$B = (\rho_n(r_s)A_n + \rho_p(r_s)A_p)/\rho(r_s), \quad (59)$$

where $f = 1 - \nu/(m+1)$. For non-trivial solutions to exist in the core, we must have $\det W = 0$, which leads to $f = 0$ or $f = \rho_{12}(\rho_n + \rho_p)/(\rho_n\rho_p)$. For the case of $f = 0$ (i.e. $\kappa_0 = 2/(m+1)$), solutions may be given by

$$A_n = A_p = B. \quad (60)$$

This solution obviously corresponds to the r^o -mode since the oscillation frequency κ_0 is exactly the same as that of the r^o -mode, and the two superfluids flow together in the core. For the case of $f = \rho_{12}(\rho_n + \rho_p)/(\rho_n\rho_p)$ (i.e., $\kappa_0 \neq 2/(m+1)$), on the other hand, solutions are given by

$$(\rho_n(r_s)A_n + \rho_p(r_s)A_p)/\rho(r_s) = B = 0. \quad (61)$$

This solution corresponds to the r^s -modes found by Lee & Yoshida (2002), for which the two superfluids counter-move in the core (see, also, Andersson & Comer 2001). It is to be noted that the solution (60) for arbitrary η and the solution (61) for $\eta = 0$ can be derived even without the assumption that the densities ρ_n , ρ_p , and ρ_{12} do not depend on the radial distance r .

6 CONCLUSION

We have discussed the modal properties of inertial modes of rotating neutron stars with the core filled with neutron and proton superfluids, taking account of entrainment effects between the superfluids. We find that the inertial modes are split into two families, ordinary fluid inertial modes (i^o -mode) and superfluid inertial modes (i^s -mode). The two superfluids in the core counter-move for the i^s -modes. For the i^o -modes, the dimensionless frequency κ_0 is only weakly dependent on the entrainment parameter η . For the i^s -modes, on the other hand, $|\kappa_0|$ almost linearly increases as the entrainment parameter η is increased. Because of the different dependences of κ_0 on the parameter η , avoided crossings as functions of η are common between i^o - and i^s -modes. We also find that some of the i^s -modes that are unstable against the gravitational radiation reaction at $\eta = 0$ become stable invalidating the instability criterion when η becomes larger than some critical η_{crit} , the value of which

depends on the mode. The radiation driven instability associated with the current multipole radiation is quite weak for the inertial modes. Since the mutual friction damping in the superfluid core is quite strong for a typical value of η compared with other damping mechanisms considered in this paper, we conclude that the inertial modes are easily damped by the mutual friction.

Recently, Arras et al. (2002) considered non-linear couplings between an unstable r -mode and stable inertial modes of rotating normal fluid neutron stars as a mechanism that limits the amplitude growth of the r -mode, and discussed in the WKB limit the modal properties of the inertial modes, which are important to determine the saturation amplitude of the r -mode. In their analysis, however, no effects of buoyancy and possible core superfluidity on the inertial modes are correctly included, which substantially affect the modal properties such as the frequency spectra and damping rates (see Yoshida & Lee 2000b). If we consider cold neutron stars in a neutron star binary, inertial modes in the superfluid core of the stars will be excited by the tidal interaction with the companion, and the mutual friction dissipation produced by the excited inertial modes will play an important role in the energy and angular momentum transfer between the orbital motion and the neutron stars, since no g -modes are propagating in the superfluid core (Lee 1995, Andersson & Comer 2001). In fact, the damping timescale of the inertial modes propagating in the superfluid core may be given by $\tau \sim \tau_{MF}^0/\bar{\Omega}$ sec, and it is for $\bar{\Omega} \sim 0.1$ of order of 0.1 to 1 sec, which is extremely short compared with the damping timescales of g -modes due to viscous processes and emission of gravitational waves (see, e.g., McDermott, Van Horn, & Hansen 1988; Reisenegger & Goldreich 1992). Considering that the orbital period decrease $d \ln P/dt$ due to gravitational radiation reactions is proportional to D^{-4} (e.g., Shapiro & Teukolsky 1983) but that due to the tidal interaction to D^{-5} (e.g., Savonije & Papaloizou 1983) where D is the binary separation, the tidal interaction in terms of the inertial modes in the superfluid core becomes important as the binary separation decreases.

Acknowledgments: S.Y. acknowledges financial support from Fundação para a Ciência e a Tecnologia (FCT) through project SAPIENS 36280/99.

APPENDIX A: OSCILLATION EQUATIONS

For the oscillation equations in the superfluid core, we employ vectors $\mathbf{s}^n$, $\mathbf{s}^p$, $\mathbf{t}^n$, $\mathbf{t}^p$, $\mathbf{h}^n$, and $\mathbf{h}^p$, whose components are given by

$$s_k^n = S_l^n, \quad s_k^p = S_{l'}^p, \quad t_k^n = T_{l'}^n, \quad t_k^p = T_{l'}^p, \quad h_k^n = H_l^n, \quad h_k^p = H_{l'}^p, \quad (\text{A1})$$

where $l = |m| + 2(k-1)$, $l' = l+1$ for even modes and $l = |m| + 2k-1$, $l' = l-1$ for odd modes, where $k = 1, 2, 3, \dots$. Using these vectors, the perturbed continuity equations (1) and (2) are written as

$$r \frac{d\mathbf{s}^n}{dr} = - \left(\frac{d \ln \rho_n}{d \ln r} + 3 \right) \mathbf{s}^n + \Lambda_0 \mathbf{h}^n, \quad (\text{A2})$$

$$r \frac{d\mathbf{s}^p}{dr} = - \left(\frac{d \ln \rho_n}{d \ln r} + 3 \right) \mathbf{s}^p + \Lambda_0 \mathbf{h}^p. \quad (\text{A3})$$

The independent components of equations (26) and (27) are reduced to

$$\begin{aligned} & L_0 r \frac{d\mathbf{h}^n}{dr} - M_1 r \frac{d\mathbf{t}^n}{dr} - m\nu \Lambda_0^{-1} r \frac{d\mathbf{s}^n}{dr} + \{(2+m\nu)\mathbf{1} - 2m\nu \Lambda_0^{-1}\} \mathbf{h}^n - (\mathbf{1} + 2m\nu \Lambda_0^{-1}) \mathbf{s}^p \\ & + \nu \Lambda_0^{-1} C(\Lambda_1 - 2\mathbf{1}) \mathbf{t}^n + \frac{\rho_{12}}{\rho_n} \left\{ r \frac{d}{dr} (\mathbf{h}^p - \mathbf{h}^n) + 2(\mathbf{h}^p - \mathbf{h}^n) - (\mathbf{s}^p - \mathbf{s}^n) \right\} \\ & + r \frac{d}{dr} \left(\frac{\rho_{12}}{\rho_n} \right) (\mathbf{h}^p - \mathbf{h}^n) = 0, \end{aligned} \quad (\text{A4})$$

$$\begin{aligned} & L_0 r \frac{d\mathbf{h}^p}{dr} - M_1 r \frac{d\mathbf{t}^p}{dr} - m\nu \Lambda_0^{-1} r \frac{d\mathbf{s}^p}{dr} + \{(2+m\nu)\mathbf{1} - 2m\nu \Lambda_0^{-1}\} \mathbf{h}^p - (\mathbf{1} + 2m\nu \Lambda_0^{-1}) \mathbf{s}^p \\ & + \nu \Lambda_0^{-1} C(\Lambda_1 - 2\mathbf{1}) \mathbf{t}^p + \frac{\rho_{12}}{\rho_p} \left(1 + \frac{m_e}{m_p} \right)^{-1} \left\{ r \frac{d}{dr} (\mathbf{h}^n - \mathbf{h}^p) + 2(\mathbf{h}^n - \mathbf{h}^p) - (\mathbf{s}^n - \mathbf{s}^p) \right\} \\ & + r \frac{d}{dr} \left\{ \frac{\rho_{12}}{\rho_p} \left(1 + \frac{m_e}{m_p} \right)^{-1} \right\} (\mathbf{h}^n - \mathbf{h}^p) = 0, \end{aligned} \quad (\text{A5})$$

$$L_1 \mathbf{t}^n - M_0 \mathbf{h}^n + \nu K \mathbf{s}^n + \frac{\rho_{12}}{\rho_n} (\mathbf{t}^p - \mathbf{t}^n) = 0, \quad (\text{A6})$$

$$L_1 \mathbf{t}^p - M_0 \mathbf{h}^p + \nu K \mathbf{s}^p + \frac{\rho_{12}}{\rho_p} \left(1 + \frac{m_e}{m_p} \right)^{-1} (\mathbf{t}^n - \mathbf{t}^p) = 0, \quad (\text{A7})$$

where $\nu = 2\Omega/\sigma = 2/\kappa_0$ and $\mathbf{1}$ stands for a unit matrix. Here, the quantities, L_0 , L_1 , C , K , Λ_0 , Λ_1 , M_0 , and M_1 are matrices, given as follows:

For even modes,

$$\begin{aligned} (L_0)_{i,i} &= 1 - \frac{m\nu}{l(l+1)}, & (L_1)_{i,i} &= 1 - \frac{m\nu}{(l+1)(l+2)}, \\ (C)_{i,i} &= -(l+2)J_{l+1}^m, & (C)_{i+1,i} &= (l+1)J_{l+2}^m, \\ (K)_{i,i} &= \frac{J_{l+1}^m}{l+1}, & (K)_{i,i+1} &= -\frac{J_{l+2}^m}{l+2}, \\ (\Lambda_0)_{i,i} &= l(l+1), & (\Lambda_1)_{i,i} &= (l+1)(l+2), \\ (M_0)_{i,i} &= \nu \frac{l}{l+1} J_{l+1}^m, & (M_0)_{i,i+1} &= \nu \frac{l+3}{l+2} J_{l+2}^m, \\ (M_1)_{i,i} &= \nu \frac{l+2}{l+1} J_{l+1}^m, & (M_1)_{i+1,i} &= \nu \frac{l+1}{l+2} J_{l+2}^m, \end{aligned}$$

where $l = |m| + 2i - 2$ for $i = 1, 2, 3, \dots$, and

$$J_l^m \equiv \left[\frac{(l+m)(l-m)}{(2l-1)(2l+1)} \right]^{1/2}. \quad (\text{A8})$$

For odd modes,

$$\begin{aligned} (L_0)_{i,i} &= 1 - \frac{m\nu}{l(l+1)}, & (L_1)_{i,i} &= 1 - \frac{m\nu}{l(l-1)}, \\ (C)_{i,i} &= (l-1)J_l^m, & (C)_{i,i+1} &= -(l+2)J_{l+1}^m, \\ (K)_{i,i} &= -\frac{J_l^m}{l}, & (K)_{i+1,i} &= \frac{J_{l+1}^m}{l+1}, \\ (\Lambda_0)_{i,i} &= l(l+1), & (\Lambda_1)_{i,i} &= l(l-1), \\ (M_0)_{i,i} &= \nu \frac{l+1}{l} J_l^m, & (M_0)_{i+1,i} &= \nu \frac{l}{l+1} J_{l+1}^m, \\ (M_1)_{i,i} &= \nu \frac{l-1}{l} J_l^m, & (M_1)_{i,i+1} &= \nu \frac{l+2}{l+1} J_{l+1}^m, \end{aligned}$$

where $l = |m| + 2i - 1$ for $i = 1, 2, 3, \dots$

The oscillation equations in the superfluid core which we use are given as a set of linear ordinary differential equations for the variables $\mathbf{s}^X$ and $\mathbf{t}^X$ with $X = n, p$, which are obtained by eliminating the vectors $\mathbf{h}^X$ in equations (A2)–(A5) using the algebraic equations (A6) and (A7). The oscillation equations in the normal fluid envelope can be obtained by substituting $\mathbf{s}^n = \mathbf{s}^p = \mathbf{s}^o$, $\mathbf{t}^n = \mathbf{t}^p = \mathbf{t}^o$ and $\rho_n = \rho_p = \rho$ into the oscillation equations in the superfluid core, where $\mathbf{s}^o$ and $\mathbf{t}^o$ means the fluid velocity fields in the normal fluid envelope.

REFERENCES

- Akmal, A., Pandharipande, V. R., & Ravenhall, D. G. 1998, Phys. Rev. C, 58, 1804
 Alpar, M. A., Langer, S. A., & Sauls, J. A. 1984, ApJ, 282, 533
 Andersson, N. 1998, ApJ, 502, 708
 Andersson, N., & Comer, G. L. 2001, MNRAS, 328, 1129
 Andersson, N., Comer, G. L., & Langlois, D. 2002, Phys. Rev. D, 66, 104002
 Andersson, N., & Kokkotas, K. D. 2001, Int. J. Mod. Phys. D, 10, 381
 Andersson, N., Kokkotas, K. D., & Schutz, B. F. 1999, ApJ, 510, 846
 Andersson, N., Kokkotas, K. D., & Stergioulas, N. 1999, ApJ, 516, 307
 Arras, P., Flanagan, E. E., Morsink, S. M., Schenk, A. K., Teukolsky, S. A., & Wasserman, I. 2003, ApJ, in press (astro-ph/0202345).
 Baym, G., & Pethick, C. 1979, ARA&A, 17, 415
 Bildsten, L. 1998, ApJ, 501, L89
 Borumand, M., Joynt, R., & Kluźniak, W. 1996, Phys. Rev. C, 54, 2745
 Comer, G. L. 2002, Found. Phys., 32, 1903
 Comer, G. L., Langlois, D., & Lin, L. M. 1999, Phys. Rev. D, 60, 104025
 Cutler, L., & Lindblom, L. 1987, ApJ, 314, 234
 Epstein, R. I. 1988, ApJ, 333, 880
 Flowers, E., & Itoh, N. 1979, ApJ, 230, 847
 Friedman, J. L., & Morsink, S. M. 1998, ApJ, 502, 714
 Friedman, J. L., & Schutz, B. F. 1978, ApJ, 222, 281

- Lee, U. 1995, *A&A*, 303, 515
- Lee, U., & Saio, H. 1986, *MNRAS*, 221, 365
- Lee, U., & Yoshida, S. 2003, *ApJ*, 586, 403
- Lindblom, L., & Ipser, J. R. 1999, *Phys. Rev. D*, 59, 044009
- Lindblom, L., & Mendell, G. 1994, *ApJ*, 421, 689
- Lindblom, L., & Mendell, G. 2000, *Phys. Rev. D*, 61, 104003
- Lindblom, L., Owen, B. J., & Morsink, S. M. 1998, *Phys. Rev. Lett*, 80, 4843
- Lockitch, K. H., & Friedman, J. L. 1999, *ApJ*, 521, 764
- McDermott, P. N., Van Horn, H. M., & Hansen, C. J. 1988, *ApJ*, 325, 725
- Mendell, G. 1991a, *ApJ*, 380, 515
- Mendell, G. 1991b, *ApJ*, 380, 530
- Owen, B. J., Lindblom, L., Cutler, C., Schutz, B. F., Vecchio, A., Andersson, N. 1998, *Phys. Rev. D*, 58, 084020
- Prix, R., & Rieutord, M. 2002, *A&A*, 393, 949
- Reisenegger, A., & Goldreich, P. 1992, *ApJ*, 395, 240
- Savonije, G. J., & Papaloizou, J. C. B. 1983, *MNRAS*, 203, 581
- Sedrakian, A., & Wasserman, I. 2000, *Phys. Rev. D*, 63, 024016
- Shapiro, S. L., & Teukolsky, S. A. 1983, *Black Holes, White Dwarfs, and Neutron Stars* (New York: Wiley)
- Thorne, K. 1980, *Rev. Mod. Phys.*, 52, 299
- Unno, W., Osaki, Y., Ando, H., Saio, H., & Shibahashi, H. 1989, *Nonradial Oscillations of Stars* (2d ed.; Tokyo: Univ. Tokyo Press)
- Yoshida, S., & Lee, U. 2000a, *ApJ*, 529, 997
- Yoshida, S., & Lee, U. 2000b, *ApJS*, 129, 353
- Yoshida, S., & Lee, U. 2003, *Phys. Rev. D*, in press (gr-qc/0304073)

Table 1. κ_0 for the i^o - and i^s -modes for the $\eta = 0$ model

$l_0 - m $	i^o -mode	i^s -mode	i -mode
1	-	-	0.6667
2	1.0978	1.1187	1.1000
	-0.5685	-0.5115	-0.5566
3	1.3564	1.3865	1.3578
	0.5180	0.5078	0.5173
	-1.0293	-0.9729	-1.0259
4	1.5191	1.5522	1.5196
	0.8639	0.8611	0.8630
	-0.2738	-0.2421	-0.2753
	-1.2738	-1.2242	-1.2729

Table 2. κ_0 and τ_i^0 's for the i^o - and i^s -modes of $m = 2$

Modes	$l_0 - m $	η	κ_0	$\tau_{GJ,2}^0$ (s)	$\tau_{GJ,3}^0$ (s)	$\tau_{GJ,4}^0$ (s)	$\tau_{GJ,5}^0$ (s)	τ_S^0 (s)	τ_{MF}^0 (s)
i^o	2	0	1.0978		-4.1×10^4		-2.4×10^{14}	2.5×10^6	∞
i^s			1.1187		-2.5×10^5		-1.2×10^{15}	1.2×10^6	∞
i^o	3		1.3564	-2.2×10^8		-1.7×10^9		3.3×10^6	∞
i^s			1.3865	-3.7×10^9		-3.5×10^{10}		5.2×10^5	∞
i^o			0.5180	-4.5×10^6		-2.1×10^6		2.6×10^6	∞
i^s			0.5078	-5.9×10^7		-1.8×10^7		5.7×10^5	∞
i^o	4		1.5191		-2.1×10^{11}		-2.4×10^{14}	3.8×10^6	∞
i^s			1.5522		-5.9×10^{12}		-1.5×10^{16}	2.2×10^5	∞
i^o			0.8639		-1.8×10^7		-1.5×10^{10}	9.9×10^6	∞
i^s			0.8611		-2.0×10^8		-5.0×10^{11}	2.3×10^6	∞
i^o	2	0.02	1.0912		-1.5×10^7		-2.1×10^{15}	6.5×10^5	9.1×10^{-1}
i^s			1.4333		-1.8×10^9		-9.5×10^{14}	6.1×10^5	1.8×10^{-1}
i^o	3		1.3591	-2.1×10^8		-1.7×10^9		2.8×10^7	3.8×10^1
i^s			1.7829	-1.2×10^{13}		-1.4×10^{17}		1.6×10^5	1.9×10^{-1}
i^o			0.5171	-4.1×10^6		-1.9×10^6		8.3×10^6	1.4×10^1
i^s			0.6464	-3.2×10^8		-1.8×10^{11}		3.0×10^5	2.4×10^{-1}
i^o	4		1.5239		-1.4×10^{13}		-2.7×10^{14}	1.6×10^7	3.3×10^0
i^s			1.9832		-8.9×10^{22}		-2.3×10^{34}	1.8×10^5	1.3×10^{-1}
i^o			0.8640		-1.7×10^7		-1.5×10^{10}	2.3×10^7	1.4×10^2
i^s			1.1063		-1.2×10^6		-8.9×10^{14}	1.8×10^5	2.3×10^{-1}
i^o	2	0.04	1.1015		-3.5×10^4		-4.9×10^{14}	2.6×10^7	2.2×10^1
i^s			1.7632		-4.1×10^{12}		-8.4×10^{19}	4.7×10^5	4.9×10^{-2}
i^o	3		1.3605	-2.2×10^8		-1.7×10^9		3.0×10^7	7.0×10^0
i^s			2.1661	1.2×10^{14}		5.9×10^{18}		1.3×10^5	4.5×10^{-1}
i^o			0.5172	-4.4×10^6		-1.9×10^6		4.1×10^6	4.2×10^0
i^s			0.7831	-3.0×10^8		-2.4×10^{10}		1.9×10^5	6.7×10^{-2}
i^o	4		1.5188		-9.8×10^{10}		-2.4×10^{14}	4.0×10^6	3.3×10^0
i^s			2.4083		2.2×10^{13}		8.9×10^{19}	6.3×10^3	3.7×10^{-2}
i^o			0.8642		-1.9×10^7		-1.5×10^{10}	2.2×10^7	3.9×10^1
i^s			1.3476		-1.2×10^{10}		-2.2×10^{15}	1.1×10^5	5.3×10^{-2}




Figure 1. First three expansion coefficients S_l^v and T_l^v for the i^o -mode with $m = 2$ and $l_0 - |m| = 2$ for the case of $\eta = 0$, given as a function of r/R . The mode has $\kappa_0 = -0.5686$. The amplitudes are normalized by $T_{m+1}^v = 1$ at $r = R$.




Figure 2. Same as Figure 1 but for S_l^w and T_l^w .




Figure 3. First three expansion coefficients S_l^v and T_l^v for the i^s -mode with $m = 2$ and $l_0 - |m| = 2$ for the case of $\eta = 0$, given as a function of r/R . The mode has $\kappa_0 = -0.5115$. The amplitudes are normalized by $T_{m+1}^v = 1$ at $r = R$.




Figure 4. Same as Figure 3 but for S_l^w and T_l^w .




Figure 5. κ_0 's for even parity inertial modes having positive κ_0 , given as a function of η . The labels $i^X(k)$ signify the i^X -mode of $l_0 - |m| = k$, where X is “o” or “s”.




Figure 6. Same as Figure 5 but for even parity inertial modes having negative κ_0 .




Figure 7. κ_0 's for odd parity inertial modes having positive κ_0 , given as a function of η . The labels $i^X(k)$ signify the i^X -mode of $l_0 - |m| = k$, where X is “o” or “s”. Here, $r^s(1)$ indicates the superfluid r -mode, which is the i^s -mode of $l_0 - |m| = 1$.




Figure 8. Same as Figure 7 but for odd parity inertial modes having negative κ_0 .




Figure 9. First three expansion coefficients S_l^v and T_l^v for the i^o -mode with $m = 2$ and $l_0 - |m| = 2$ for the case of $\eta = 0.01$, given as a function of r/R . The mode has $\kappa_0 = -0.5403$. The amplitudes are normalized by $T_{m+1}^v = 1$ at $r = R$.




Figure 10. Same as Figure 9 but for S_l^w and T_l^w .




Figure 11. First three expansion coefficients S_l^v and T_l^v for the i^s -mode with $m = 2$ and $l_0 - |m| = 2$ for the case of $\eta = 0.01$, given as a function of r/R . The mode has $\kappa_0 = -0.6103$. The amplitudes are normalized by $T_{m+1}^v = 1$ at $r = R$.




Figure 12. Same as Figure 11 but for S_l^w and T_l^w .